

Vacuum energy, spectral determinant and heat kernel asymptotics of graph Laplacians with general vertex matching conditions

J. M. Harrison* and K. Kirsten†

*Department of Mathematics, Baylor University,
Waco, TX 76798, USA*

**E-mail: jon.harrison@baylor.edu*

†E-mail: klaus.kirsten@baylor.edu

We consider Laplace operators on metric graphs, networks of one-dimensional line segments (bonds), with matching conditions at the vertices that make the operator self-adjoint. Such quantum graphs provide a simple model of quantum mechanics in a classically chaotic system with multiple scales corresponding to the lengths of the bonds. For graph Laplacians we briefly report results for the spectral determinant, vacuum energy and heat kernel asymptotics of general graphs in terms of the vertex matching conditions.

1. Introduction

Quantum graphs provide a simple, analytically tractable, model in which to investigate quantum phenomena. This is of particular interest for the study of vacuum energy and related spectral quantities as the spectrum of a quantum graph generically incorporates the features associated with quantum mechanical systems that are classically chaotic.¹ In particular the eigenvalues of the Laplace operator on a graph are typically distributed as the eigenvalues of large random matrices where the appropriate random matrix ensemble for comparison is determined by the symmetries of the quantum system; this correspondence is known as the Bohigas-Giannoni-Schmit conjecture.²

Quantum graphs were introduced as a model of quantum chaos by Kotlos and Smilansky.³ Graph models are also of current interest in many areas of mesoscopic physics like Anderson localization, photonic crystals, microelectronics, nanotechnology and the theory of wave guides; see Ref. 4 for a review. In this article we present results for the vacuum energy,^{5,6} spectral determinant⁷⁻⁹ and heat kernel asymptotics^{10,11} of graph Laplacians. The

unified approach we adopt, based on the spectral zeta function of the graph, provides new results in each case. The full description of the construction of zeta functions of quantum graphs will appear in Ref. 12.

2. Graph Laplacian

Let $G = (\mathcal{V}, \mathcal{B})$ be a graph where $\mathcal{V}$ is the set of vertices and $\mathcal{B}$ is the set of bonds (or edges); for an example see Fig. 1. Each bond $b \in \mathcal{B}$ joins a pair of vertices and for notational convenience we assume that bonds are directed with an initial and terminal vertex specified via the functions $o : \mathcal{B} \rightarrow \mathcal{V}$ and $t : \mathcal{B} \rightarrow \mathcal{V}$ respectively. We insist that the set $\mathcal{B}$ is symmetric so that $b \in \mathcal{B}$ if and only if there is another bond $\bar{b} \in \mathcal{B}$ the reversal of b such that $o(b) = t(\bar{b})$ and $t(b) = o(\bar{b})$. The total number of bonds $B = |\mathcal{B}|$ is therefore twice the physical number of connections in the network. Each bond b is associated with an interval $[0, L_b]$ so that the local coordinate $x_b = 0$ at $o(b)$ and $x_b = L_b$ at $t(b)$. We naturally require that $L_{\bar{b}} = L_b$ so there is a single length associated with a physical connection in the network, consequently $x_{\bar{b}} = L_b - x_b$. The total length of the graph G is denoted by $\mathcal{L} = \frac{1}{2} \sum_{b=1}^B L_b$.




Fig. 1. A star (or hydra) graph.

A function ψ on G is defined by the collection of functions $\{\psi_b(x_b)\}_{b \in \mathcal{B}}$ on the bonds of the graph. We require $\psi_b(x_b) = \psi_{\bar{b}}(L_b - x_b)$ in order that ψ be physically meaningful. The Hilbert space on the graph is

$$L^2(G) = \bigoplus_{b=1}^B L^2([0, L_b]) . \quad (1)$$

To define a Laplacian on G we consider the operator $-\frac{d^2}{dx_b^2}$ on the bonds with matching conditions between functions ψ_b meeting at vertices of G to make the graph Laplacian self-adjoint: for example Neumann like matching conditions require ψ to be continuous at the vertices and $\sum_{b|o(b)=v} \psi'_b(0) =$

0 for all vertices $v \in \mathcal{V}$. All such self-adjoint matching conditions were classified by Kostykin and Schrader¹³ and we follow their scheme. The matching conditions on the whole graph are specified by the pair of $B \times B$ matrices $\mathbb{A}, \mathbb{B}$ via

$$\mathbb{A}\psi + \mathbb{B}\psi' = \mathbf{0} , \quad (2)$$

where $\psi = (\psi_1(0), \dots, \psi_B(0))^T$ and $\psi' = (\psi'_1(0), \dots, \psi'_B(0))^T$. These matching conditions define the domain of a self-adjoint Laplace operator iff $\text{rank}(\mathbb{A}, \mathbb{B}) = B$ and $\mathbb{A}\mathbb{B}^\dagger = \mathbb{B}\mathbb{A}^\dagger$.

To find the graph spectrum we want to solve the Laplace eq. on the bonds

$$-\frac{d^2}{dx_b^2}\psi_b(x_b) = k^2\psi_b(x_b) . \quad (3)$$

In the case of a star graph, Fig. 1, solving Eq. (3) on the bonds and substituting in the matching condition at the center one obtains the well known secular eq. of the star graph $\sum_{b=1}^B \tan kL_b = 0$, whose solutions $\{k_j | k_j \geq 0\}$ are square roots of the eigenvalues of the Laplacian on the star.

If instead one substitutes solutions of (3) in the general form of matching conditions (2) we find the sequence $\{k_j\}$ are the solutions of a secular eq. $f(k) = 0$ where,

$$f(k) = \det \left(\mathbb{A} + k\mathbb{B} \begin{pmatrix} -\cot(k\mathbf{L}) & \csc(k\mathbf{L}) \\ \csc(k\mathbf{L}) & -\cot(k\mathbf{L}) \end{pmatrix} \right) , \quad (4)$$

and $\mathbf{L} = \text{diag}\{L_1, \dots, L_B\}$. This can be compared to the scattering matrix formulation introduced by Kottos and Smilansky.^{3,14}

3. Results

The zeta function of the quantum graph can be formulated as a contour integral applying the argument principle to (4) following the technique introduced in Refs. 15,16, namely

$$\zeta(s) = \sum_{j=0}^{\infty} ' k_j^{-2s} = \frac{1}{2\pi i} \int_c z^{-2s} \frac{f'(z)}{f(z)} dz = \frac{1}{2\pi i} \int_c z^{-2s} \frac{d}{dz} \log f(z) dz . \quad (5)$$

The prime on the sum denotes the exclusion of zero modes if present. Generically the poles of f are the whole of the set $\{m\pi/L_b | m \in \mathbb{Z}, b = 1, \dots, B\}$ and this is the case we consider.^a The contour c is then chosen so as to

^aIt is possible for particular choices of matching conditions $\mathbb{A}, \mathbb{B}$ and bond lengths $\{L_b\}$ that individual poles cancel in the determinant, however, a small perturbation of the bond lengths removes this.

include the zeros of f on the positive real axis and exclude the poles. Making a contour transformation to an integral along the imaginary axis closed with a semicircular arc on the right we obtain an expression for $\zeta(s)$ valid in the strip $0 < \Re s < 1$,

$$\zeta(s) = \zeta_R(2s) \sum_{b=1}^B \left(\frac{\pi}{L_b} \right)^{-2s} + \frac{\sin \pi s}{\pi} \int_0^\infty t^{-2s} \frac{d}{dt} \log \hat{f}(t) dt, \quad (6)$$

$$\hat{f}(t) = \det \left(\mathbb{A} - t\mathbb{B} \begin{pmatrix} \coth(t\mathbf{L}) & -\operatorname{csch}(t\mathbf{L}) \\ -\operatorname{csch}(t\mathbf{L}) & \coth(t\mathbf{L}) \end{pmatrix} \right), \quad (7)$$

where ζ_R is the Riemann zeta function.

Regularized formulations of the vacuum energy, and the spectral determinant as well as the heat kernel asymptotics are obtained directly from the zeta function. The vacuum energy, formally $E = \frac{1}{2} \sum_{j=0}^\infty 'k_j^2$ where k_j^2 is an eigenvalue of the Laplacian, is defined as $E_c = \frac{1}{2} \zeta(-1/2)$. Differentiating with respect to the L_β , the length of the bond β , we obtain the Casimir force on the bond,

$$F_c^\beta = \frac{\pi}{24L_\beta^2} + \frac{1}{\pi} \int_0^\infty \frac{\partial}{\partial L_\beta} \log \hat{f}(t) dt. \quad (8)$$

The spectral determinant is formally the product of the eigenvalues $\prod_{j=0}^\infty 'k_j^2$. The zeta function representation (6) allows a direct evaluation of the regularized spectral determinant, as

$$\det'(-\Delta) = \exp(-\zeta'(0)) = \frac{2^B \hat{f}(0)}{c_N \prod_{b=1}^B L_b}. \quad (9)$$

Here, c_N denotes the first non-zero coefficient in the $t \rightarrow \infty$ asymptotic expansion of

$$\hat{f}(t) \sim \det(\mathbb{A} - t\mathbb{B}) = \det \mathbb{B} t^{2B} + c_{2B-1} t^{2B-1} + \cdots + c_1 t + \det \mathbb{A}. \quad (10)$$

So $c_N = \det \mathbb{B}$ when $\det \mathbb{B} \neq 0$. The factor of c_N in (9) is introduced by the analytic continuation of $\zeta(s)$ to $s = 0$.

The heat kernel for a quasi one-dimensional system has an expansion for $t \rightarrow 0$ of the form, $K(t) = \sum_{j=1}^\infty e^{-k_j^2 t} \sim \sum_{\ell=0,1/2,1,\dots}^\infty a_\ell t^{\ell-1/2}$. The heat kernel coefficients are related to the zeta function¹⁷ by $a_\ell = \operatorname{Res}(\zeta(s)\Gamma(s))|_{s=1/2-\ell}$. Subtracting sufficiently many terms in the $t \rightarrow \infty$ expansion of $\hat{f}$, the zeta function (6) can be extended as far left of the imaginary axis as required and we obtain the full asymptotic expansion of

the heat kernel of the general graph Laplacian,

$$K(t) \sim \frac{\mathcal{L}}{\sqrt{4\pi t}} + \frac{N}{2} - \sum_{k=1,3/2,2,\dots}^{\infty} \frac{b_{2k-1}}{\Gamma(k - \frac{1}{2})} t^{k-1/2} . \quad (11)$$

The coefficients b_n are obtained from the expansion of $\log \hat{f}(t)$ at infinity,

$$\log \hat{f}(t) \sim N \log t + \log c_N + \sum_{n=1}^{\infty} \frac{b_n}{t^n} . \quad (12)$$

For a given set of matching conditions defined by matrices $\mathbb{A}, \mathbb{B}$ the coefficients b_n can be easily obtained with a computer algebra package.

Acknowledgments

The authors would like to thank G Berkolaiko, JP Keating, P Kuchment, J Marklof, R Piziak and B Winn for helpful suggestions. KK is supported by National Science Foundation grant PHY-0554849 and JMH is supported by National Science Foundation grant DMS-0604859.

References

1. S. Gnuzmann and U. Smilansky, *Adv. Phys.* **55**, 527–625 (2006).
2. O. Bohigas, M.-J. Giannoni and C. Schmit, *Phys. Rev. Lett.* **52**, 1-4 (1984).
3. T. Kottos and U. Smilansky, *Phys. Rev. Lett.* **79**, 4794–4797 (1997).
4. P. Kuchment, *Waves Random Media* **12**, R1–24 (2002).
5. S. Fulling, L. Kaplan and J. H. Wilson, *Phys. Rev. A* **76**, 012118 (2007).
6. G. Berkolaiko, J. M. Harrison and J. H. Wilson, *J. Phys. A: Math. Theor.* **42**, 025204 (2009).
7. J. Desbois, *J. Phys. A: Math. Gen.* **33**, L63–L67 (2000).
8. J. Desbois, *Eur. Phys. J. B* **24**, 261–266 (2001).
9. L. Friedlander, in: G. Berkolaiko, R. Carlson, S. A. Fulling, and P. Kuchment (Eds) *Quantum Graphs and Their Applications*, Contemporary Mathematics, **415**, 151–160 (AMS, 2006).
10. V. Kostykin, J. Potthoff and R. Schrader, in F. Germinet and P. D. Hislop (Eds), *Adventures in mathematics*, Contemporary Mathematics **447**, 175–198 (AMS, 2007).
11. J. Bolte and S. Endres, *Ann. Henri Poincaré* **10**, 189–223 (2009).
12. J. M. Harrison and K. Kirsten, “Zeta functions of quantum graphs,” (Preprint) [arXiv:0911.2509](https://arxiv.org/abs/0911.2509).
13. V. Kostykin and R. Schrader, *J. Phys. A: Math. Gen.* **32**, 595–630 (1999).
14. T. Kottos and U. Smilansky, *Ann. Phys.* **274**, 76–124 (1999).
15. K. Kirsten and A. J. McKane, *Ann. Phys.* **308**, 502–527 (2003).
16. K. Kirsten and A. J. McKane, *J. Phys. A: Math. Gen.* **37**, 4649–4670 (2004).
17. R.T. Seeley, in *Singular Integrals*, Chicago. Proc. Sympos. Pure Math. **10**, 288–307 (1966), (AMS, 1968).